

Seminar
CITTA' IN SALUTE
Thursday 15 October
h. 14.00-14.30
Amber Room 1

Green areas and public health: improving wellbeing and physical activity in the urban context

Spazi verdi e salute pubblica: migliorare il benessere e l'attività fisica nei contesti urbani

Daniela D'Alessandro,¹ Maddalena Buffoli,² Lorenzo Capasso,³ Gaetano Maria Fara,⁴ Andrea Rebecchi,² Stefano Capolongo,^{2,5} and the Hygiene on Built Environment Working Group on Healthy Buildings of the Italian Society of Hygiene, Preventive Medicine and Public Health (SIItI)

¹Dip. Ingegneria civile, edile e ambientale, Sapienza Università di Roma, Italy; ²Dip. Architettura, ingegneria delle costruzioni e ambiente costruito, Politecnico di Milano, Italy; ³Dip. Sanità pubblica, medicina sperimentale e forense, Università di Pavia, Italy; ⁴Dip. Sanità pubblica e malattie infettive, Sapienza Università di Roma, Italy; ⁵coordinator of the Hygiene on Built Environment WG

Corresponding author: Stefano Capolongo; e-mail: stefano.capolongo@polimi.it

Abstract

Introduction. The green areas play an important role in improving the environmental quality and climate of the city. However, despite the undoubted importance of these areas, the benefits to public health are still under investigation.

Results. The Authors perform a review of studies on the relationship between health and green areas and describe the main areas for which evidence on this relationship is currently available. They include: effects on air quality, on social cohesion, on mental health, with particular reference to the stress, and on physical activity.

Conclusions. Most of the evidence comes from cross-sectional and descriptive studies and the approaches used to measure the association show themselves to be often unsatisfactory. The Authors conclude the review stressing the need for greater integration between the different professionals involved in urban planning and in health care analysis in order to identify research approaches more appropriate to understand such complex issues, striving toward a planning design of green areas that will satisfy both environmental sustainability and health requirements.

(*Epidemiol Prev* 2015; 39(4) Suppl 1: 8-13)

Key words: green space, urban context, health, physical activity, design

Riassunto

Introduzione. Le aree verdi assumono un ruolo molto importante nel migliorare la qualità ambientale e il clima delle città. Tuttavia, nonostante l'indiscussa importanza di tali aree, i benefici per la salute pubblica sono ancora oggetto di studio.

Risultati. E' stata svolta una revisione degli studi sul rapporto tra aree verdi e salute. Sono descritti i principali ambiti sui quali sono oggi disponibili evidenze circa la relazione tra spazi verdi e salute. Essi includono: gli effetti sulla qualità dell'aria, sulla coesione sociale, sulla salute mentale, con particolare riferimento allo stress e sull'attività fisica.

Conclusioni. La maggior parte delle evidenze derivano da studi descrittivi e di prevalenza e gli approcci utilizzati per la misura dell'associazione sono spesso non soddisfacenti. Gli autori concludono la revisione sottolineando la necessità di una maggiore integrazione tra le diverse professionalità coinvolte nella progettazione urbana e nella tutela della salute per individuare approcci di ricerca più adeguati per comprendere problematiche così complesse, in modo da giungere a una progettazione delle aree verdi che soddisfi le esigenze di sostenibilità ambientale e di salute.

(*Epidemiol Prev* 2015; 39(4) Suppl 1: 8-13)

Parole chiave: spazi verdi, contesti urbani, salute, attività fisica, progettazione

INTRODUCTION

About 3.4 billion people (half of the world's population) live in urban areas today: it has been estimated that the proportion will rise to 60% by 2030, and the absolute number will be 6.3 billion by 2050.¹ The understanding of how urban environment affects health outcomes and how it could produce health benefits is therefore an urgent priority, as recognized by WHO in the declaration of 2010 as the Year of Urban Health.

Unfortunately, due to increasing urbanization, combined with the spatial planning policies of densification, nowadays homes have become more and more separated from green environments, although, as argued by van de Berg et al.,² a restricted access to green spaces may increase vulnerability to the impact of stressful life events on mental and physical health.

At present, several studies provide evidence, albeit still rather weak, of some association between green spaces, wellbeing, and health.³⁻⁶ In the most recent years, this topic has gained an increasing interest.

RESULTS

A «web of knowledge» search with just two terms, «green space and health», yielded 2 hits for 1990-1999, 34 for 2000-2009, and 45 from 2010 to June 2013.⁷ The reasons for this rise in interest are many. Some papers reflect concerns that urbanization, environment degradation, and lifestyle changes are quali-quantitatively diminishing occasions for human contact with nature; others consider nature as just one aspect of the physical environment that may be potentially beneficial for health. In general, research aimed to propose practical measures that could directly or indirectly provide access to nature, including interventions about housing, transportation, and recreation.

In the same paper,⁷ the Authors performed a «review of reviews» on the topic until April 2013, involving 56 relevant reviews. Of the peer-reviewed articles, the majority appeared in journals focused on public health or environmental planning.

The reviews varied considerably in methodology, guiding aims, and environmental aspects discussed.


Despite this variety in methods, aims, and focus, there was a strong agreement about the methodological state of the art. The reviews agreed about the following conclusions:

- most of the studies are observational;
- few primary studies have investigated in a consistent and rigorous way the relationships between contact with nature and health;
- few primary studies have explored how the effects may vary by population subgroup, type of natural environment, or type of contact with nature;
- consistent and objective measurements of both exposure to nature and health-related outcomes remain elusive.

Reviews showed three common weaknesses:

- search strategies did not always address the variety of environments or settings that count as “natural”;
- reviews from individual-based disciplines tended to ignore evidence from others (e.g., psychology *vs.* epidemiology);
- the variety of health outcomes used in primary studies was not well addressed.

Nevertheless, the reviews generally agreed that beneficial effects from contact with nature do occur. Contact with nature may affect health via multiple pathways, that have received relatively large amounts of research attention: air quality, social cohesion, stress reduction, and physical activity (figure 1). Pathways mostly proceed from the natural environment to health effects via contact with nature, but two of them also proceed directly from the natural environment to stress and air quality, respectively, implying that the natural environment may affect air quality and health without engaging first with nature as such. The two-headed arrow between “natural environment” and “contact with nature as such” acknowledges that these are connected in a two-way relationship. The two-headed arrows between the boxes describing air quality, physical activity, social contacts, and stress also imply their recip-



rocal relatedness; however, each may be related with all the others and not only with the ones closeby.

Air pollution

Interactions between green spaces and health may lead to positive and negative effects, involving both social and physical health effects.⁸ Impermeable surfaces such as roofs, facades, and roads impair the microclimate by disturbing the radiation and energy balance of areas. *Vice versa*, green areas in one's living environment may ameliorate air pollution, and the urban heat island effect.^{9,10} Trees, shrubs, and other kind of vegetation may affect air quality and, through it, human wellbeing and health. Trees and other vegetation may reduce levels of some pollutants, including gases and particulate matter (PM), but they may also contribute to air pollution by releasing hydrocarbons, including isoprene and terpenes, with considerable variation by species.¹⁰⁻¹² Some trees and plants release pollens, aggravating asthma and allergies. In fact urban and sub-urban ecological changes can affect the geographical range of diseases related to pollens, insects, and parasites.^{13,14} Finally, trees improve air quality indirectly when they cool urban environments and reduce building energy demand.⁷ Hence, the positive benefits of green space cannot be generalized.

Social cohesion

Natural features and open spaces in a residential area can play an important role in residents' feelings of attachment towards the community, and their interactions with other residents. Social cohesion refers to shared norms and values, the existence of positive and friendly relationships, and feelings of being accepted and of enjoying membership. It is more a characteristic of neighbourhoods than of individuals¹⁵ and therefore more likely to be influenced by physical characteristics of the neighborhood, such as the availability and quality of green spaces and natural elements. At the same time, green spaces that are perceived to have become overgrown or to be unmanaged, may have a negative effect on peoples' wellbeing by increasing anxiety caused by fear of crime.⁷ Kneeshaw-Price et al.¹⁶ evaluated five neighbourhood crime-related safety measures to determine how they were interrelated. They found that neighbourhood-active children living in the lowest crime-quartile neighborhoods had 40 minutes more of total MVPA (moderate to vigorous physical activity) in comparison to neighborhood-active children living in the highest crime-quartile neighbourhoods.

Generally, the few available studies suggest a positive relationship between social cohesion and natural environments.^{7,17} As argued by Hartig et al.,⁷ social cohesion within a neighborhood does not lend itself to experimental research, which makes it difficult to determine whether relations with environmental features are causal. Some researchers¹⁸ found a positive association between the presence of trees and grass in common spaces and informal social contact with neighbours. Moreover, the relationship between greenery and social contacts appeared to be mediated by the use of the common spaces. The research also showed that social contact was positively related to one's sense

of safety. Subsequent studies showed that residents with more trees and grass around their buildings displayed less aggressive behaviour, and their buildings were associated with fewer crimes.^{19,20} More recent research suggests similar positive effects of greening vacant lots, especially a reduction in gun assaults and disorderly conduct.²¹

Stressful life

In general, individuals living in areas that lack green spaces may be more vulnerable to the negative impacts of stressful life events because they have less opportunities for nature-based coping strategies than individuals living in areas with abundant green spaces.² Thus, the availability of green space in the surroundings may be an important environmental factor that moderates the relationship between stressful life events and health. According to the WHO, mental health promotion should include actions that create living conditions and environments that support mental health and allow people to adopt and maintain healthy lifestyles.²² Green spaces are thought to influence mental health through an increase in physical activity, by providing places for neighbourhood residents to meet, encouraging social ties, and alleviating stress and mental fatigue.²³ Van de Berg et al.² observed that respondents with a higher amount of green space in a 3 km radius were less affected by experiencing a stressful life and had a better perceived mental health. The moderating effects of green space were found only for those within 3 km, presumably because the 3 km indicator is more affected by the presence of larger areas of green space.

Nevertheless, findings from the studies on green spaces and mental health relationship are not homogeneous. Most studies were cross-sectional, and are thus subject to reverse causality. Gascon et al.,²⁴ in a recent systematic review, found limited evidence of a causal relationship between surrounding greenness and mental health in adults, whereas the evidence was inadequate in children. However, at least two longitudinal studies in the UK provided evidence that individuals living in greener areas had better mental health outcomes over time,^{25,26} while a study in Sweden found an additive protective effect of green space and physical activity on mental health among women.²⁷ Astell-Burt et al.²⁸ reported a protective effect of green space on minor psychiatric morbidity across the lifecourse in early adulthood for men. In contrast, the benefit of green space for women emerged later in adulthood.

Despite the advantages of longitudinal designs, concerns about unmeasured confounds remain, most notably the inability to control for non-random selection of residents into neighborhoods. For this reason Cohen-Cline et al.²³ examined the association between access to green space and mental health among adult twin pairs, a way to address this self-selection problem because they provide a method of controlling genetic and environmental confounds. Authors have hypothesized that greater access to green space is associated with less depression, but provide less evidence for effects on stress or anxiety. Access to green spaces showed psychological health benefits for socioeconomically deprived populations, in at least three stud-

ies;²⁹⁻³¹ green space may therefore narrow health inequalities. A recent large, cross-sectional international study,³² although not proving causality, concludes that socioeconomic inequalities in mental wellbeing were smaller among urban dwellers reporting good access to recreational/green areas and there was no such difference for the other tested neighborhood services. There are reasons that support the plausibility of a causal effect: for example, green spaces between residences and heavily trafficked roads can reduce occupant noise annoyance, vegetation can conceal displeasing structures, and landscaping around housing can help residents maintain privacy and avoid feelings of crowding.^{33,34}

Hordyk et al.,³⁵ investigating everyday practices of immigrant children and families in the context of urban green spaces such as parks, fields, backyards, streetscapes, gardens, forests and rivers, observed that activities in natural environment serve as a protective factor in the health and wellbeing of this population, providing emotional and physical nourishment in the face of adversity. Participants accessed urban nature to minimize the effects of inadequate housing, strengthen social cohesion and reduce emotional stress.

Physical activity

A growing body of evidence suggests that a major determinant of physical activity is access to green spaces. In 2004, the *Guide for Community Preventive Services* of the CDC recommended «creation of or enhanced access to places for physical activity based on strong evidence of their effectiveness in increasing physical activity and improving physical fitness».^{36,37} As the *Guide* noted, most of the evidence for this recommendation derived from cross-sectional studies. Low rates of physical activity are an important contributing factor to rising levels of obesity, as well as risk of cardiovascular disease, hypertension, type 2 diabetes, stroke, colon cancer, and premature death. Physical activity also promotes mental health across the life span.^{38,39} Recent evidence suggests that the health benefits of increased physical activity are largest among those who were initially physically inactive⁴⁰ and in youth.⁴¹⁻⁴³ The outdoor environment may influence how much physically active an individual is by offering suitable spaces for certain types of performances. It may also attract people outdoors because of the experiences it offers. Such outings ordinarily entail some form of physical activity, at least walking. An important precondition for the use of natural environments for physical activity is individuals' (perceived) safety.^{7,16} Although there are possible negative effects associated with physical activity (e.g., sports injuries) as well as with being in a natural environment (e.g., Lyme disease from tick bites), most of these effects are not specific to physical activity in a natural environment. The built environment is an important factor of influence, as it can facilitate or inhibit participation in physical activity. The literature has examined how different aspects of public open space, such as access to, size, and design features, are associated with participation in physical activity. Proximity to parks and recreational settings are generally associated with greater physical activity.⁴⁴ Qualitative evidence further shows that safety,

aesthetics, amenities, maintenance, and proximity of public open spaces are important attributes for supporting physical activity.⁴⁴

Despite the increasing number of studies in this field, there are some inconsistencies that confuse urban designers and policy makers and prevent the development of evidence-based guidelines. Koohsari et al.⁴⁴ in a recent review identified conceptual and methodological gaps that need to be addressed to progress research on public open space and physical activity, which include:

- **uneven definition of open space**, which introduces difficulties in comparing and collating evidence across different studies. While there is a lack of research into the influence different types of public open space have on physical activity, there is some evidence that non-park public open spaces might be important for physical activity (e.g., walking trails). As such, there might be specific requirements for designing a walking trail to accommodate a wide range of physical activities within a small linear place compared with a park.
 - **use of longitudinal and experimental study design when possible**: in particular research on public open spaces and physical activity would benefit from experimental studies that measure behaviours before and after the introduction of new public open space or renovation of existing public open space. This point could take advantage from the increase in urban regeneration interventions that several countries, including Italy, are promoting. A recent study of people who relocated from one neighbourhood to another, found that gaining access to three different types of public open space (parks, sport fields, beach) increased daily walking by 18-20 minutes for each type of public open space gained.⁴⁵
 - **exploring public open space exposure in multiple contexts**: previous studies have primarily focused on public open space in a residential context, while the effect of public open space in other settings (e.g., around workplaces and schools) has been ignored. Karusisi et al.⁴⁶ found that the number of supermarkets around workplaces was associated with walking for transport among workers. Dalton et al.⁴⁷ observed that active travel to work was negatively associated with the availability of free car parking at workplaces.⁴⁸
 - **moving specific measures of physical activity into public open space**: public open space can influence activity in at least three ways:
 - a. it can be a setting where people engage in physical activity;
 - b. it can be a destination to which people actively travel either to be active or simply to socialize;
 - c. it can be used as part of a route to pass through to reach another destination. Hence public open space can contribute to different types of physical activity behaviours. Few studies have attempted to understand the variety of ways public open spaces influence physical activity.
- Koohsari et al.⁴⁴ also considered the importance of the quality of public open space, in terms of features and characteristics related to physical activity or walking, but also on the char-

acteristics of surrounding built environment and on the association between public open space type and user profiles (age group, socio-demographic group, sex). Furthermore they stress the need to identify threshold values needed to attract people to public open space.

CONCLUSIONS

Providing adequate public green spaces means offering people both physical and psychological health benefits. As argued by Wolch et al.,¹⁰ most studies reveal that the distribution of such space often disproportionately benefits the most affluent communities. Giving everybody access to green spaces is therefore increasingly recognized as an issue of environmental justice. Many cities in other countries (e.g., both the US and China) have implemented strategies to increase the supply of urban green space, especially in deprived neighborhoods. Strategies include greening of marginal urban land and reuse of obsolete or underutilized infrastructures.

The definition of appropriate and innovative solutions could benefit from collaboration between different professions.^{49,50} Therefore it is necessary to encourage the integration of information between various professional figures, such as urban na-

ture conservationists, urban planners, environmental psychologists, and public health specialists.^{49,51} Hartig et al.⁷ underline that a lot more remains to be done to help environmental policy makers and designers to establish realistic assessments of what nature can and cannot do in their domain of activity. Part of this task involves explaining how health benefits might overlap (or conflict) with other benefits, such as better storm water management, species preservation, and carbon sequestration. At the same time, a comprehensive methodology for analysis of the associations between aspects of the urban environment and residents' health needs to be identified and implemented.⁵² As argued by Rydin et al.,¹ the absence of such a methodology is largely attributable to the complex nature of urban systems, in which many factors affect social and health outcomes, compounded by the scarcity of consistent data available at the urban scale. In this field, Koohsari et al.⁴⁴ suggest several indications to improve research approaches. Building a body of evidence in this way could contribute to provide much-needed data to urban designers to plan a public open space system able to promote public health.

Conflicts of interest: none declared

References/Bibliografia

- Rydin Y, Bleahu A, Davis M, et al. Shaping cities for health: complexity and the planning of urban environments in the 21st century. *Lancet* 2012;379:2079-108.
- van de Berg AE, Maas J, Verheij RA, Groenewegen PP. Green space as a buffer between stressful life events and health. *Soc Sci Med* 2010;70:1203-1210.
- Tzoulas K, Korpela K, Venn S, et al. Promoting ecosystem and human health in urban areas using green infrastructure: a literature review. *Landscape Urban Plann* 2007;81:167-78.
- Bai H, Wilhelm Stanis SA, Kaczynski AT, Besenyi GM. Perceptions of neighborhood park quality: associations with physical activity and body mass index. *Ann Behav Med* 2013;45:S39-S48.
- Pearson AL, Bentham G, Day P, Kingham S. Associations between neighbourhood environmental characteristics and obesity and related behaviors among adult New Zealanders. *BMC Public Health* 2014;14:553.
- Takano T, Nakamura K, Watanabe M. Urban residential environments and senior citizens' longevity in mega-city areas: the importance of walkable green space. *J Epidemiol Commun Health* 2002;56:913-16.
- Hartig T, Mitchell R, de Vries S, Frumkin H. Nature and Health. *Annu Rev Public Health* 2014;35:207-28.
- Signorelli C, Riccò M. The health environment interaction in Italy. *Ig Sanità Pubbl* 2012;68:374-80. PMID: 23074745.
- Whitford V, Ennos AR, Handley JF. City form and natural processes: indicators for the ecological performance of urban areas and their application to Merseyside, UK. *Landscape Urban Plann* 2001;20:91-103.
- Wolch JR, Byrne J, Newell JP. Urban green space, public health, and environmental justice: the challenge of making cities "just green enough". *Landscape Urban Plann* 2014;125:234-44.
- Benjamin MT, Winer AM. Estimating the ozone-forming potential of urban trees and shrubs. *Atmos Environ* 1998;32:53-68.
- Buffoli M, Capolongo S, Loconte VL, Signorelli C. Thermovalorization: new technologies, impacts and mitigation strategies. *Ann Ig* 2012;24:167-78. PMID: 22755503.
- Patz JA, Norris DE. Land use change and human health. *Ecosyst Land Use Change* 2004;153:159-67.
- Zielinski-Gutierrez EC, Hayden MH. A model for defining West Nile Virus risk perception based on ecology and proximity. *EcoHealth* 2006;3:28-34.
- Baum FE, Ziersch AM, Zhang G, Osborne K. Do perceived neighborhood cohesion and safety contribute to neighbourhood differences in health? *Health Place* 2009;15:925-34.
- Kneeshaw-Price SH, Saelens BE, Sallis JF, et al. Neighborhood crime-related safety and its relation to children's physical activity. *J Urban Health* 2015;3:472-89.
- de Vries S, van Dillen SME, Groenewegen PP, Spreeuwenberg P. Streetscape greenery and health: stress, social cohesion and physical activity as mediators. *Soc Sci Med* 2013;94:26-33.
- Kuo FE, Sullivan WC, Coley RL, Brunson L. Fertile ground for community: inner-city neighborhood common spaces. *Am J Community Psychol* 1998;26: 823-51.
- Kuo FE, Sullivan WC. Aggression and violence in the inner city: effects of environment via mental fatigue. *Environ Behav* 2001;33:543-71.
- Kuo FE, Sullivan WC. Environment and crime in the inner city: Does vegetation reduce crime? *Environ Behav* 2001;33:343-67.
- Branas CC, Cheney RA, MacDonald JM, Tam VW, Jackson TD, Ten Have TR. A difference-indifferences analysis of health, safety, and greening vacant urban space. *Am J Epidemiol* 2011;174:1296-306.
- WHO Mental Health: *Strengthening Our Response*. <http://www.who.int/mediacentre/factsheets/fs220/en/> (accessed June 2, 2015).
- Cohen-Cline H, Turkheimer E, Duncan GE. Access to green space, physical activity and mental health: a twin study. *J Epidemiol Community Health* 2015; 69:523-29.
- Gascon M, Triguero-Mas M, Martinez D, et al. Mental health benefit of long-term exposure to residential green and blue spaces: a systematic review. *Int J Environ Res Public Health* 2015;12:4354-79.
- Alcock I, White MP, Wheeler BW, et al. Longitudinal effects on mental health of moving to greener and less green urban areas. *Environ Sci Technol* 2014; 48:1247-55.
- White MP, Alcock I, Wheeler BW, et al. Would you be happier living in a greener urban area? A fixed-effects analysis of panel data. *Psychol Sci* 2013;24:920-28.
- Annerstedt M, Ostergren PO, Bjork J, et al. Green qualities in the neighbourhood and mental health – Results from a longitudinal cohort study in Southern Sweden. *BMC Public Health* 2012;12:337.
- Astell-Burt T, Mitchell R, Hartig T. The association between green space and mental health varies across the lifecycle. A longitudinal study. *J Epidemiol Community Health* 2014;68:578-83.

29. Maas J, Verheij RA, Groenewegen PP, de Vries S, Spreeuwenberg P. Green space, urbanity, and health: how strong is the relation? *J Epidemiol Community Health* 2006;60:587-92.
30. Lachowycz K, Jones AP. Does walking explain associations between access to greenspace and lower mortality? *Soc Sci Med* 2014;107:9-17.
31. Mitchell R, Popham F. Effect of exposure to natural environment on health inequalities: an observational population study. *Lancet* 2008;372:1655-60.
32. Mitchell RJ, Richardson EA, Shortt NK, Pearce JR. Neighborhood environments and socioeconomic inequalities in mental well-being. *Am J Prev Med* 2015. In press.
33. Capolongo S, Buffoli M, Oppio A, Petronio M. Sustainability and Hygiene of building: future perspectives. *Epidemiol Prev* 2014;38(Suppl 2):46-50. Italian.
34. Capolongo S, Buffoli M, Oppio A, Rizzitello S. Measuring hygiene and health performance of buildings: a multidimensional approach. *Ann Ig* 2013;25:151-57. PMID: 23471453.
35. Hordyk SR, Hanley J, Richard E. "Nature is there; its free": Urban greenspace and the social determinants of health of immigrant families. *Health Place* 2015; 6:74-82.
36. Centers for Disease Control and Prevention. *Environmental and Policy Approaches to Increase Physical Activity: Creation of or Enhanced Access to Places for Physical Activity Combined with Informational Outreach Activities 2004*. <http://www.thecommunityguide.org/pa/environmentalpolicy/improvingaccess.html> (accessed June 2, 2015).
37. Heath GW, Brownson RC, Kruger J, Miles R, Powell KE, Ramsey LT, and the Task Force for Community Preventive Services. The effectiveness of urban design and land use and transportation policies and practices to increase physical activity: a systematic review. *J Phys Act Health* 2006;3:S55-S76.
38. Bize R, Johnson JA, Plotnikoff RC. Physical activity level and health-related quality of life in the general adult population: a systematic review. *Prev Med* 2007;45:401-15.
39. Janssen I, LeBlanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act* 2010;7:40.
40. Powell KE, Paluch AE, Blair SN. Physical activity for health: What Kind? How much? How Intense? On Top of What? *Annu Rev Public Health* 2011;32: 349-65.
41. Jann I, Rosu A. Undeveloped green space and free-time physical activity in 11 to 13-year-old children. *Int J Behav Nutr Phys Act* 2015;12:26.
42. Sanders T, Feng X, Fahey PP, Lonsdale C, Astell-Burt T. Green neighbourhoods, slimmer children? Evidence from 4,423 participants aged 6 to 13 years in the Longitudinal Study of Australian children. *Int J Obesity* 2015. In press.
43. Christian H, Zubrick SR, Foster S, et al. The influence of the neighborhood physical environment on early child health and development: a review and call for research. *Health & Place* 2015;33:25-36.
44. Koohsari MJ, Mavoa S, Villanueva K, et al. Public open space, physical activity, urban design and public health: concepts, methods and research agenda. *Health & Place* 2015;33:75-82.
45. Giles-Corti B, Bull F, Knuiam M, et al. The influence of urban design on neighbourhood walking following residential relocation: longitudinal results from the RESIDE study. *Soc Sci Med* 2013;77:20-30.
46. Karusisi N, Thomas F, Meline J, Brondeel R, Chaix B. Environmental conditions around itineraries to destinations as correlates of walking for transportation among adults. The RECORD cohort study. *Plos One* 2014;9:e88929.
47. Dalton AM, Jones AP, Panter JR, Ogilvie D. Neighborhood, route and work-place-related environmental characteristics predict adults' mode of travel to work. *Plos One* 2013;8(6): e67575.
48. Buffoli M, Capolongo S, Bottero M, Cavagliato E, Speranza S, Volpatti L. Sustainable Healthcare: how to assess and improve healthcare structures' sustainability. *Ann Ig* 2013;25:411-18. PMID: 24048179.
49. D'Alessandro D, Raffo M. Adeguare le risposte ai nuovi problemi dell'abitare in una società che cambia. *Ann Ig* 2011;23:267-74.
50. Capolongo S, Buffoli M, Oppio A. How to assess the effects of urban plans on environment and health. *Territorio* 2015;73:145-51.
51. Capolongo S, Battistella A, Buffoli M, Oppio A. Healthy design for sustainable communities. *Ann Ig* 2011;1:43-53.
52. Capasso L, Capolongo S, Flacco ME, Manzoli L, Buffoli M. Housing quality and its health impact in Italy: current knowledge, methodological issues, new evaluation models. *Ig Sanita Pubbl* 2014 Jul Aug;70(4):411-22. Italian. PMID: 25353271.